

12 児童生徒の疾病異常（令和4年度・公立）

（保健体育課調）

| 区 分 | | 小学校 | | 中学校 | | 義務教育学校 (前期) | | 義務教育学校 (後期) | | 高等学校 (全日制) | |
|------------|-------------------|-------|-------|-------|-------|----------------|-------|----------------|-------|---------------|-------|
| | | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 |
| 裸眼視力 | 1.0未満0.7以上の者 | 11.8% | 13.1% | 14.0% | 14.9% | 8.6% | 7.3% | 9.8% | 14.8% | 16.1% | 16.8% |
| | 0.7未満0.3以上の者 | 11.7% | 13.4% | 18.1% | 19.6% | 6.6% | 10.6% | 18.0% | 17.8% | 19.8% | 19.1% |
| | 0.3未満の者 | 5.9% | 7.1% | 12.9% | 15.4% | 6.8% | 7.7% | 12.8% | 15.2% | 15.9% | 13.7% |
| 眼 | 感染性眼疾患の者 | 0.0% | 0.0% | 0.0% | 0.0% | 0% | 0% | 0% | 0% | 0% | 0% |
| | その他の眼疾患・異常の者 | 5.4% | 4.6% | 5.6% | 4.6% | 5.5% | 3.7% | 3.4% | 5.4% | 3.8% | 3.2% |
| 聴 | 難聴の者 | 1.0% | 1.2% | 0.2% | 0.4% | 0% | 0.5% | 0.4% | 0% | 1.1% | 1.6% |
| 耳・鼻・咽 | 耳疾患の者 | 5.7% | 5.5% | 4.6% | 3.4% | 7.6% | 4.3% | 5.1% | 3.0% | 2.5% | 1.4% |
| | 鼻・副鼻腔疾患の者 | 11.0% | 7.3% | 10.5% | 8.0% | 18.6% | 11.5% | 12.0% | 10.6% | 10.2% | 7.5% |
| | 口腔咽頭疾患の者 | 0.6% | 0.5% | 0.3% | 0.2% | 0% | 0% | 0% | 0% | 0.1% | 0.1% |
| 歯 | 処置完了の者 | 16.1% | 14.8% | 13.4% | 15.4% | 19.8% | 17.3% | 17.2% | 18.9% | 21.7% | 25.6% |
| | 未処置歯のある者 | 16.8% | 15.5% | 10.5% | 11.3% | 26.9% | 29.4% | 17.5% | 20.2% | 12.5% | 12.5% |
| | その他の疾病の者 | 6.5% | 6.7% | 4.4% | 3.8% | 14.1% | 17.3% | 7.7% | 9.4% | 2.0% | 1.9% |
| 尿 | 蛋白陽性の者 | 0.3% | 0.8% | 1.4% | 1.6% | 0.7% | 1.1% | 0.5% | 0.9% | 0.9% | 1.0% |
| | 潜血陽性の者 | 0.4% | 1.2% | 0.5% | 3.1% | 1.0% | 1.3% | 0.3% | 4.6% | 0.4% | 2.1% |
| | 糖陽性の者 | 0.1% | 0.1% | 0.1% | 0.2% | 0% | 0% | 0% | 0% | 0.2% | 0.2% |
| 寄生虫 | 寄生虫卵保有の者 | 0% | 0% | - | - | 0% | 0% | - | - | - | - |
| その他の疾病及び異常 | 栄養不良の者 | 0.0% | 0.0% | 0.1% | 0.1% | 0.5% | 0% | 0.3% | 0% | 0.0% | 0.1% |
| | 肥満傾向の者 | 1.5% | 0.9% | 1.0% | 0.5% | 3.5% | 3.0% | 0.8% | 0.9% | 0.5% | 0.2% |
| | 脊柱側弯の者 | 0.3% | 0.7% | 1.2% | 5.1% | 0.9% | 0.7% | 0.5% | 1.5% | 1.2% | 3.1% |
| | その他の脊柱・胸郭・四肢の異常の者 | 0.5% | 0.4% | 1.0% | 0.7% | 0.2% | 0% | 0.5% | 0.9% | 0.7% | 0.4% |
| | 感染性皮膚疾患の者 | 0.0% | 0.0% | 0.0% | 0% | 0% | 0% | 0% | 0% | 0.0% | 0% |
| | アトピー性皮膚炎の者 | 4.4% | 3.9% | 4.1% | 3.6% | 4.5% | 2.6% | 3.6% | 0.9% | 2.5% | 1.9% |
| | 心臓疾患・異常の者 | 1.0% | 1.0% | 1.1% | 1.0% | 0.5% | 1.5% | 0.5% | 0.3% | 0.9% | 0.9% |
| | ぜん息の者 | 5.7% | 3.9% | 4.5% | 3.0% | 8.4% | 6.2% | 4.1% | 1.8% | 2.5% | 2.0% |
| | 腎臓疾患の者 | 0.3% | 0.4% | 0.4% | 0.5% | 0.5% | 0.7% | 0% | 0% | 0.3% | 0.2% |
| 結核 | 精密検査を受けた者 | 0.2% | 0.2% | 0.1% | 0.1% | 0% | 0.2% | 0% | 0% | 0.1% | 0.2% |
| | 結核の者 | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% |

注・定期健康診断結果による。

・「0」は皆無、「0.0」は数値が表章単位未満、「-」は実施しない場合である。