

マイワシの粗脂肪量

水揚日: 令和2年4月28日


船名: 33 不動丸 漁場: 36°02'N 140°55'E (鹿島沖)

Lipid content of Japanese sardines

Landing day : April 28, 2020

Fishing ground : Off the coast of Kashima, Japan

36°02'N 140°55'E



No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content	No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content
1	14.9	37.4	4.7	11	17.0	50.6	5.9
2	21.1	107.6	15.8	12	15.2	36.3	5.9
3	17.6	53.3	5.9	13	20.7	86.4	10.3
4	16.2	43.6	3.5	14	17.2	59.5	8.4
5	18.8	74.1	7.2	15	17.0	50.9	7.2
6	15.3	38.0	11.5	16	15.7	36.8	7.2
7	17.7	60.0	10.9	17	22.0	118.3	16.4
8	15.4	36.2	5.9	18	14.9	32.8	5.9
9	20.4	104.8	25.1	19	17.7	58.7	9.0
10	19.0	74.2	10.3	20	22.9	132.6	24.5

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議