

マイワシの粗脂肪量

水揚日: 令和2年2月14日


船名: 53 波一丸 漁場: 36°35'N 140°59'E(日立沖)

Lipid content of Japanese sardines

Landing day : February 14, 2020

Fishing ground : Off the coast of Hitachi, Japan

36°35'N 140°59'E



No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content	No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content
1	19.7	106.4	17.1	11	19.3	97.9	18.3
2	19.4	90.8	14.0	12	19.8	92.0	12.7
3	20.2	115.4	20.1	13	19.9	82.8	9.6
4	20.1	106.8	15.8	14	19.3	97.6	22.6
5	18.8	100.5	25.1	15	19.3	85.0	9.6
6	18.6	79.1	17.7	16	17.3	81.6	20.8
7	19.7	101.5	23.2	17	17.9	67.9	9.6
8	18.5	72.1	9.0	18	16.7	60.9	11.5
9	19.1	100.4	14.6	19	16.6	59.5	14.6
10	19.9	91.6	12.7	20	14.3	33.4	1.0

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議