

マイワシの粗脂肪量

水揚日: 令和2年2月3日


船名: 57 角石丸 漁場: 36°39'N 141°02'E (日立沖)

Lipid content of Japanese sardines

Landing day : February 3, 2020

Fishing ground : Off the coast of Hitachi, Japan

36°39'N 141°02'E



No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content	No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content
1	19.5	112.7	26.3	11	17.1	60.2	17.1
2	20.5	102.5	14.6	12	14.8	44.2	14.0
3	20.6	114.9	20.8	13	16.2	52.6	15.2
4	21.2	102.5	12.1	14	18.4	76.7	10.3
5	20.0	132.6	30.0	15	16.8	67.9	22.0
6	20.0	96.3	14.6	16	16.7	56.6	12.1
7	19.2	74.7	14.0	17	17.5	76.1	25.1
8	17.3	63.0	12.7	18	14.6	38.9	14.0
9	16.3	56.4	13.4	19	15.5	42.7	10.9
10	18.2	67.8	10.9	20	16.2	58.2	17.1

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議