

マイワシの粗脂肪量

水揚日: 令和2年5月8日


船名: 35 不動丸 漁場: 35°45'N 141°05'E (犬吠埼沖)

Lipid content of Japanese sardines

Landing day : May 8, 2020

Fishing ground : Off the coast of Inubosaki, Japan

35°45'N 141°05'E



No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content	No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content
1	23.1	141.9	2.2	11	18.9	84.7	1.6
2	22.7	138.0	12.7	12	19.0	84.7	3.5
3	22.3	145.0	15.8	13	18.8	82.8	4.1
4	21.9	126.9	4.1	14	18.8	79.1	7.8
5	22.7	137.8	1.6	15	18.7	76.9	5.3
6	22.2	124.5	6.6	16	17.6	70.0	4.1
7	19.4	85.8	12.1	17	16.7	58.4	4.7
8	19.1	82.5	2.9	18	17.9	62.6	1.6
9	19.6	90.2	0.4	19	17.9	70.0	4.7
10	21.2	105.5	10.3	20	18.6	79.2	4.1

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議