

マサバの粗脂肪量

水揚日: 令和元年12月10日

船名: 33不動丸 漁場: 37°08'N 141°28'E (福島沖)

Lipid content of chub mackerels

Landing day : December 10, 2019

Fishing ground : Off the coast of Fukushima, Japan

37°08'N 141°28'E



No.	尾叉長cm Fork length	体重g Body weight	粗脂肪量% Lipid content	No.	尾叉長cm Fork length	体重g Body weight	粗脂肪量% Lipid content
1	34.2	553.9	23.1	11	33.6	502.0	23.1
2	34.9	554.9	24.1	12	34.1	439.7	17.0
3	35.3	593.1	25.1	13	33.1	461.5	19.0
4	36.1	613.6	22.6	14	31.8	389.3	21.0
5	33.0	462.8	25.6	15	34.1	516.5	24.1
6	33.6	449.3	21.5	16	32.5	395.1	18.0
7	34.2	502.0	22.6	17	31.2	321.9	10.9
8	33.1	444.6	18.0	18	30.6	342.3	20.0
9	35.0	474.8	17.0	19	31.7	371.6	22.1
10	32.5	470.0	25.6	20	31.1	337.9	15.0

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議