

マイワシの粗脂肪量

水揚日:令和2年1月14日


船名:8 成田不動丸 漁場:36°10'N 140°51'E(鹿島沖)

Lipid content of Japanese sardines

Landing day : January 14, 2020

Fishing ground : Off the coast of Kashima, Japan

36°10'N 140°51'E



No.	体長cm	体重g	粗脂肪量%	No.	体長cm	体重g	粗脂肪量%
	Standard length	Body weight	Lipid content		Standard length	Body weight	Lipid content
1	23.5	152.0	17.7	11	18.0	67.7	19.5
2	21.6	136.8	30.0	12	18.2	64.0	2.2
3	21.6	111.3	14.0	13	19.1	71.8	15.8
4	20.7	127.2	28.2	14	17.8	61.5	16.4
5	19.9	106.1	22.0	15	17.6	59.2	12.1
6	19.1	92.4	24.5	16	16.6	50.0	17.1
7	20.0	103.0	20.8	17	16.4	50.3	20.1
8	20.1	90.6	20.8	18	16.6	59.7	20.1
9	20.5	114.8	25.1	19	16.3	44.8	8.4
10	19.6	76.5	8.4	20	16.5	50.5	13.4

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議